

# OSTEOLOGICAL EVALUATION

*Prepared by*  
**EVAN MATSHES BSc, MD**  
*Consultant Osteologist*



Product No. BC-217

**Human Male European  
Skull, Hammer Blows**



**Bone Clones, Inc.**

OSTEOLOGICAL REPRODUCTIONS

9200 Eton Ave. Chatsworth, CA 91311

Phone: (818) 709-7991 or (800) 914-0091 (USA only)

Email: [info@boneclones.com](mailto:info@boneclones.com) Web: [www.boneclones.com](http://www.boneclones.com)

© Bone Clones, Inc. 2015

# **Human, Male, White, Healed hammer blows**

**Product Number:** BC-217

**Specimen Evaluated:** Bone Clones® replica

**Skeletal Inventory:** 1 intact cranium

## **General observations:**

In general, the molding process has preserved significant details necessary for evaluation. The general shape and configuration of the skull is within normal limits. The ectocranial morphology of the individual cranial bones is within normal limits. The sutural patterns are of expected configuration; however, there is advanced fusion/ossification. There is the suggestion of bilateral epipteric bones. The foramina are of expected configuration.

## **Dentition:**

There are 10 teeth in the maxillary arcade. All teeth have an adult morphology and no deciduous dentition remains. The dentition is atraumatic. There are no dental restorations or prostheses. There is a moderate degree of attrition.

The following maxillary dentition is present: 1.7 [#2], 1.6 [#3], 1.3 [#6], 1.2 [#7], 1.1 [#8], 2.1 [#9], 2.3 [#11], 2.4 [#12], 2.6 [#14], and 2.7 [#15].

The atraumatic gomphoses of 1.5 [#4], 1.4 [#5], 2.2 [#10] are empty and are without signs of healing.

1.8 [#1] and 2.8 [#16], and their gomphoses, are not present.

Both 1.6 [#3] and 2.6 [#14] have cusps of Carabelli.

2.7 [#15] has a small lingual groove defect, suggestive that an amalgam filling has fallen out.

There is generalized mild to moderate periodontal disease, with the molar teeth showing furcation involvement.

**Features of Race:**

The interocular distance is not prominently widened. The nasal root is prominent and the nasal angle is acute. The zygomatic bones retreat posteriorly from the plane of the face. The nasal aperture is narrow both superiorly and inferiorly. The anterior nasal spine is short, and the inferior margin of the nasal aperture has a sharp (nasal) sill. The maxillary dental arcade has a somewhat rectangular-shape. It is not possible to assess the degree of alveolar prognathism; however, the maxillary arcade projects slightly. The maxillary incisors are blade-shaped. It is not possible to assess for edge-on-edge incisal bite. There is no post-bregmatic depression. The calvarial sutures are focally complex (at least at the extremes of the coronal suture).

***The totality of features is most in keeping with those of a White individual.***

**Features of Sex:**

There is marked prominence of the cranial sites for musculofascial attachment including especially:

- the nuchal lines
- the external occipital protuberance
- the mastoid processes of the temporal bones
- the temporal lines
- the supraorbital tori
- the occipital condyles
- the supramastoidal crest

The nasion is somewhat rough, and the supraorbital margins are blunted.

***The totality of features is most in keeping with male sex.***

## ***Bone Clones® Osteological Evaluation Report***

### **Features of Age:**

There are no identifiable fontanelles. The sphenoccipital synchondrosis is fused.

Ten ectocranial osteologic landmarks are evaluated for degree of suture closure according to the Meindl and Lovejoy method\*. [1] Scores are assigned as follows:

<b>1</b>	3
<b>2</b>	3
<b>3</b>	3
<b>4</b>	3
<b>5</b>	2
<b>6</b>	2
<b>7</b>	2
<b>8</b>	2
<b>9</b>	2
<b>10</b>	2

\* As is always the case with casting, there is a tendency towards overscoring.

The sum of scores for the cranial vault (landmarks 1 through 7) is 18. This corresponds to an estimated age of 48.8 +/- 10.5 years.

The sum of scores for the anterior cranium (landmarks 6 through 10) is 10. This corresponds to an estimated age of 51.9 +/- 12.5 years.

### **Trauma:**

A depressed circular defect is on the left side of the frontal bone, along the superior margin of the left orbit. The depression is not of uniform depth, and the margins and surfaces are smooth, indicating that it is a healed/healing injury.

A second depression, on the lateral aspect of the right parietal bone, is also vaguely circular, but has an imprinted pattern of parallel ridges. Again, the peripheral margins and surfaces of the depression are smooth, indicating that this right parietal depression is a healed/healing injury.

***The circular shape, the size, and the depressed nature of the left frontal and right parietal lesions are consistent with blunt trauma inflicted by blows with the head of a hammer.***

**SUMMARY:**

1. White.
2. Male.
3. 39.4 –59 years; range 38 – 64.4 years.
4. Two healed or healing depressed skull fractures:
  - a. Consistent with blunt trauma inflicted with a hammer.
5. No evidence of significant osteologic variations or primary pathology.

**EDUCATIONAL RESOURCES:**

1. This is an excellent example of the skull of a White male.
  - a. The concept of race assessment is controversial. It may be worthwhile to review the varying schools of thought on this issue. Short summaries from the perspective of the forensic anthropologist[2] and forensic pathologist[3] are readily available.
  - b. In many circumstances, the skull alone will allow an investigator to correctly determine sex.[4] However, the findings in the skull should never be treated in isolation; rather, they should be incorporated into your ‘whole case’ database. This database should include information obtained from all other aspects of the case. From an osteologic perspective, this includes (importantly) the bones of the pelvis.
  - c. Age assessment of skeletal remains is best done in the context of the entire skeleton. Assessment of the degree of suture closure can be used with some degree of success[1]; however, there is tremendous variability in the degree of closure process. Students must be cautioned that statistical data is based on **populations**, and may not necessarily be reflective of reality in an **individual**.
2. This is an excellent example of healed or healing blunt trauma, in a pattern that is classic for blows delivered by a hammer.
3. It may be appropriate to discuss the broad spectrum of sequelae that may develop after blunt head trauma (e.g. meningitis, intracranial hemorrhages, etc.).
4. It may be appropriate to discuss the rather critical role of radiography (including computed tomography) in the evaluation of traumatized human remains.

## ***Bone Clones*® Osteological Evaluation Report**

### **REFERENCES:**

1. Meindl, R.S. and Lovejoy, C.O. (1985). Ectocranial suture closure: a revised method for the determination of skeletal age at death based on the lateral-anterior sutures. *American Journal of Physical Anthropology*, 68(1): 57-66.
2. Gill, G. (1998). Craniofacial criteria in the skeletal attribution of race. In *Forensic Osteology: Advances in the Identification of Human Remains*, K. Reichs, Editor. Springfield, IL: Charles C. Thomas.
3. Matshes, E. and Lew, E. (2006). Forensic osteology. In *Forensic Pathology: Principles and Practice*, D. Dolinak, E. Matshes, and E. Lew, Editors. San Diego, CA: Elsevier (Academic Press).
4. Krogman, W. and Iscan, M. (1986). *The Human Skeleton in Forensic Medicine*. 2 ed. Springfield, IL: Charles C. Thomas.

### **DISCLAIMERS:**

This report is meant only as a teaching tool for introductory level students of the anatomical, anthropology or forensic sciences who might be using this specimen to learn human and forensic osteology. My opinions are based solely upon the material presented to me. This is somewhat artificial as in real forensic investigations additional studies would be undertaken prior to the formulation of diagnoses, and the production of a report. These studies might include plain film radiography, computed tomography (CT) studies, histology, etc. My opinions regarding race and sex are based only upon non-metric analyses. Evaluation of cranial suture closure is most accurately assessed endocranially as the sutures are known to close from the endocranial table towards the ectocranium. My opinions regarding this skull were made without access to the postcranial skeleton.

Evan Matshes BSc, MD  
Consultant Osteologist

Emma Lew BSc, MD  
Consultant Forensic Pathologist